

## Press release

## Living near green areas reduces the risk of suffering a stroke by 16%

- **In contrast, exposure to environmental pollutants such as fine particulate matter, nitrogen dioxide and soot significantly increases the risk of suffering an ischaemic stroke**
- **The article in the journal Environment International, one of the most important studies published to date in Europe, analysed data from the entire population in Catalonia, geo-referencing more than three and a half million people.**
- **The authors consider that these results mean that the maximum levels of atmospheric pollutants established by the EU need to be reconsidered**

**Barcelona, 15<sup>th</sup> March, 2022.** - The **risk of suffering an ischaemic stroke**, the most common type of cerebrovascular event, **is 16% less** in people who have **green spaces** less than 300 metres from their homes. This has been revealed in a joint study by the Hospital del Mar Medical Research Institute, Hospital del Mar, the Catalan Health Quality and Assessment Agency (AQuAS) from the Catalan Government's Department of Health and the Barcelona Institute for Global Health (ISGlobal), a centre promoted by the "la Caixa" Foundation. It is the most important work in this field to date in Europe, analysing data on the entire population of Catalonia between 2016 and 2017. It has been published in the journal Environment International.

The study took into account information on **exposure to three atmospheric pollutants** linked to **vehicle traffic** in more than three and a half million people selected from among the 7.5 million residents of Catalonia, over the age of eighteen who had not suffered a stroke prior to the start of the study. Specifically, it analysed the impact of the levels of particulate matter under 2.5 microns (PM<sub>2.5</sub>), nitrogen dioxide (NO<sub>2</sub>) and soot particles at the place of residence of each of the people studied. The number and density of green spaces within a 300-metre radius of their homes was also studied. The data was obtained via geographic referencing and by designing models of exposure to the two determinants, using population data obtained by reusing health information generated in Catalonia and anonymised by the Data Analytics Programme for Research and Innovation in Health (Programa d'Analítica de Dades per a la Recerca i la Innovació en Salut; PADRIS) and managed by the Catalan Agency for Health Quality and Assessment (Agència de Qualitat i Avaluació Sanitàries de Catalunya; AQuAS).

### More pollution, a higher risk of stroke

The results indicate a **direct relationship between increased levels of NO<sub>2</sub> in the atmosphere and the risk of ischaemic stroke**. For every increase of 10 micrograms (µg) per cubic metre, this risk increases by 4%. The same happens when PM<sub>2.5</sub> levels increase by 5 µg/m<sup>3</sup>. In the case of soot particles, the risk increases by 5% for every 1 µg/m<sup>3</sup> increase in the atmosphere. These figures are the same for the entire population, irrespective of other socio-economic factors, age or smoking habits.

**"It should be borne in mind that, unlike other air pollutants, which have various sources, NO<sub>2</sub> is mainly caused by road traffic. Therefore, if we really want to reduce the multiple risks that this pollutant poses to people's health, we need to implement bold measures to reduce car use"**, says Cathryn Tonne, a researcher at ISGlobal.

**"The study demonstrates the importance of environmental determinants in stroke risk. Given that it is predicted that the incidence, mortality and disability attributed to the disease will increase in the coming years, it is important to understand all the**

## Press release

**risk factors involved**", explains Carla Avellaneda, a researcher in the Neurovascular Research Group at IMIM-Hospital del Mar and one of the main authors of the study. Previous studies by the same group had already provided evidence on the relationship between factors such as soot or noise levels and the risk of suffering a stroke and its severity. All these factors act as stroke triggers.

In contrast, having an abundance of green spaces within the same radius from the home directly reduces the risk of suffering a stroke. Specifically, up to 16%. In this sense, **"People who are surrounded by greater levels of greenery at their place of residence are protected against the onset of stroke"**, says Avellaneda. Exposure to green spaces is generally considered to have beneficial effects through a variety of mechanisms, such as stress reduction, increased physical activity and social contact, and even exposure to an enriched microbiome.

### Rethinking established air pollutant limits

In view of this data, the researchers point out that it is necessary to reflect on the current levels of atmospheric pollution that are considered safe. At the moment, the thresholds set by the European Union are 40 µg/m<sup>3</sup> for NO<sub>2</sub>, which the World Health Organisation reduces to 10 µg/m<sup>3</sup>, and 25 µg/m<sup>3</sup> for PM<sub>2.5</sub>, which the WHO limits to 5 µg/m<sup>3</sup>. Currently, no levels have been set for soot particles. In fact, the levels recorded during the period analysed were lower, on average, than those set by the European authorities (17 µg/m<sup>3</sup> for PM<sub>2.5</sub>, 35 µg/m<sup>3</sup> for NO<sub>2</sub> and 2.28 µg/m<sup>3</sup> in the case of soot).

**"Despite compliance with the levels set by the European Union, we are faced with the paradox that there is still a health risk, such as the one we identified in this study, where there is a direct relationship between exposure to pollutants in our environment and the risk of suffering a stroke"**, explains Rosa Maria Vivanco, lead author of the study and researcher at AQuAS and IMIM-Hospital del Mar. **"The danger is still present and many more measures need to be taken"**, considering the increase in the urban population and its ageing, she adds.

In this sense, Jaume Roquer, head of the Neurology Service at Hospital del Mar and coordinator of the Neurovascular Research Group at IMIM-Hospital del Mar, points out that **"This study demonstrates the real impact that environmental aspects have on the health of the Catalan population. In view of the effects of atmospheric pollution, the lack of green spaces, noise, and so forth, more efforts and populational strategies are needed to reduce its impact. Its harmful effects are permanently and globally damaging. We must strive to achieve more sustainable towns and cities where living does not mean an increased risk of disease."**

### Reference article

[Air pollution and surrounding greenness in relation to ischemic stroke: A population-based cohort study](#). Environment International (IF9.621), Pub Date : 2022-02-15, DOI: [10.1016/j.envint.2022.107147](#) C. Avellaneda-Gómez, R.M. Vivanco-Hidalgo, S. Olmos, U. Lazcano, A. Valentin, C. Milà, A. Ambrós, J. Roquer, C. Tonne

### Further information

IMIM/Hospital del Mar Communications Service: Marta Calsina 93 3160680 [mcalsina@imim.es](mailto:mcalsina@imim.es), David Collantes 600402785 [dcollantes@hospitaldelmar.cat](mailto:dcollantes@hospitaldelmar.cat)

AQuAS Communications Service: [comunicacio.aquas@gencat.cat](mailto:comunicacio.aquas@gencat.cat)



Institut Hospital del Mar  
d'Investigacions Mèdiques

Salut/  Agència de Qualitat i Avaluació  
Sanitàries de Catalunya

Press release